

Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show: 20 Sort Send to Text

About Entrez

Text Version

☐ 1: Nature. 2000 Sep 14;407(6801):242-8.

Related Articles, I

nature

Vascular-specific growth factors and blood vessel formation.

Yancopoulos GD, Davis S, Gale NW, Rudge JS, Wiegand SJ, Holash J.

Regeneron Pharmaceuticals, Inc., Tarrytown, New York 10591, USA.

A recent explosion in newly discovered vascular growth factors has coincided with exploitation of powerful new genetic approaches for studying vascular development. An emerging rule is that all of these factors must be used in perfect harmony to form functional vessels. These new findings also demand re-evaluation of therapeutic efforts aimed at regulating blood vessel growth in ischaemia, cancer and other pathological settings.

Publication Types:

- Review
- Review, Tutorial

PMID: 11001067 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books
Search PubMed for AL-2 AND Eph Go Clear
Limits Preview/Index History Clipboard Details

No items found.

[About Entrez](#)

[Text Version](#)

Entrez PubMed

[Overview](#)
[Help](#) | [FAQ](#)
[Tutorial](#)
[New/Noteworthy](#)
[E-Utilities](#)

PubMed Services

[Journals Database](#)
[MeSH Database](#)
[Single Citation Matcher](#)
[Batch Citation Matcher](#)
[Clinical Queries](#)
[LinkOut](#)
[Cubby](#)

Related Resources

[Order Documents](#)
[NLM Catalog](#)
[NLM Gateway](#)
[TOXNET](#)
[Consumer Health](#)
[Clinical Alerts](#)
[ClinicalTrials.gov](#)
[PubMed Central](#)


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books
Search PubMed for AL-2 AND ephrin Go Clear
Limits Preview/Index History Clipboard Details

No items found.

[About Entrez](#)

[Text Version](#)

Entrez PubMed

[Overview](#)
[Help | FAQ](#)
[Tutorial](#)
[New/Noteworthy](#)
[E-Utilities](#)

PubMed Services

[Journals Database](#)
[MeSH Database](#)
[Single Citation Matcher](#)
[Batch Citation Matcher](#)
[Clinical Queries](#)
[LinkOut](#)
[Cubby](#)

Related Resources

[Order Documents](#)
[NLM Catalog](#)
[NLM Gateway](#)
[TOXNET](#)
[Consumer Health](#)
[Clinical Alerts](#)
[ClinicalTrials.gov](#)
[PubMed Central](#)


[Write to the Help Desk](#)

[NCBI | NLM | NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books
Search PubMed for AL-2 AND vasculature Go Clear
Limits Preview/Index History Clipboard Details

No items found.

[About Entrez](#)

[Text Version](#)

Entrez PubMed

[Overview](#)

[Help | FAQ](#)

[Tutorial](#)

[New/Noteworthy](#)

[E-Utilities](#)

PubMed Services

[Journals Database](#)

[MeSH Database](#)

[Single Citation Matcher](#)

[Batch Citation Matcher](#)

[Clinical Queries](#)

[LinkOut](#)

[Cubby](#)

Related Resources

[Order Documents](#)

[NLM Catalog](#)

[NLM Gateway](#)

[TOXNET](#)

[Consumer Health](#)

[Clinical Alerts](#)

[ClinicalTrials.gov](#)

[PubMed Central](#)


[Write to the Help Desk](#)

[NCBI | NLM | NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search **PubMed** for **EFL-6** **Go** **Clear**

Limits Preview/Index History Clipboard Details

One of your terms is not found in the database.
See [Details](#).

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Display **Summary** Show: **500** Sort **Text** Send to **Text**

Items 1 - 16 of 16

One p

☐ **1:** [Fay DS, Qiu X, Large E, Smith CP, Mango S, Johanson BL](#). [Related Articles, Li](#)

The coordinate regulation of pharyngeal development in *C. elegans* by lin-35/Rb, pha-1, and ubc-18.

Dev Biol. 2004 Jul 1;271(1):11-25.

PMID: 15196946 [PubMed - indexed for MEDLINE]

☐ **2:** [Dellaca RL, Santus P, Aliverti A, Stevenson N, Centanni S, Macklem PT, Pedotti A, Calverley PM](#). [Related Articles, Li](#)

Detection of expiratory flow limitation in COPD using the forced oscillation technique.

Eur Respir J. 2004 Feb;23(2):232-40.

PMID: 14979497 [PubMed - indexed for MEDLINE]

☐ **3:** [Baydur A, Wilkinson L, Mehdian R, Bains B, Milic-Emili J](#). [Related Articles, Li](#)

Extrathoracic expiratory flow limitation in obesity and obstructive and restrictive disorders: effects of increasing negative expiratory pressure.

Chest. 2004 Jan;125(1):98-105.

PMID: 14718427 [PubMed - indexed for MEDLINE]

☐ **4:** [Wang PH, Kuo PH, Hsu CL, Wu HD, Chang YS, Kuo SH, Yang PC](#). [Related Articles, Li](#)

Diagnostic value of negative expiratory pressure for airway hyperreactivity.

Chest. 2003 Nov;124(5):1762-7.

PMID: 14605046 [PubMed - indexed for MEDLINE]

☐ **5:** [Chen J, Laughlin LS, Hendrickx AG, Natarajan K, Overstreet JW, Lasley BL](#). [Related Articles, Li](#)

The effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on chorionic gonadotrophin activity in pregnant macaques.

Toxicology. 2003 Apr 15;186(1-2):21-31.

PMID: 12604168 [PubMed - indexed for MEDLINE]

☐ **6:** [Boxem M, van den Heuvel S](#). [Related Articles, Li](#)

C. elegans class B synthetic multivulva genes act in G(1) regulation.

Curr Biol. 2002 Jun 4;12(11):906-11.

PMID: 12062054 [PubMed - indexed for MEDLINE]


















☐ **7:** [Boni E, Corda L, Franchini D, Chiroli P, Damiani GP, Pini L, Grassi V, Tantucci C](#). [Related Articles, Li](#)

Volume effect and exertional dyspnoea after bronchodilator in patients with COPD with and without expiratory flow limitation at rest.

Thorax. 2002 Jun;57(6):528-32.

PMID: 12037229 [PubMed - indexed for MEDLINE]


☐ **8:** [Iandelli I, Aliverti A, Kayser B, Dellaca R, Cala SJ, Duranti R, Kelly S, Scano G, Sliwinski P, Yan S, Macklem PT, Pedotti A](#). [Related Articles, Li](#)

-  **Determinants of exercise performance in normal men with externally imposed expiratory flow limitation.**
J Appl Physiol. 2002 May;92(5):1943-52.
PMID: 11960944 [PubMed - indexed for MEDLINE]
-  **9: Ferretti A, Giampiccolo P, Cavalli A, Milic-Emili J, Tantucci C.** [Related Articles](#), [Li](#)
-  **Expiratory flow limitation and orthopnea in massively obese subjects.**
Chest. 2001 May;119(5):1401-8.
PMID: 11348945 [PubMed - indexed for MEDLINE]
-  **10: Michaud S, Bourgault AM, Gaudreau C.** [Related Articles](#), [Li](#)
-  **Epidemiology of enterococcal bacteremia in a referral center for hepatobiliar diseases.**
Infection. 2000 Nov-Dec;28(6):361-6.
PMID: 11139155 [PubMed - indexed for MEDLINE]
-  **11: Kumar J, van Rheeën HA.** [Related Articles](#), [Li](#)
-  **A major gene for time of flowering in chickpea.**
J Hered. 2000 Jan-Feb;91(1):67-8.
PMID: 10739130 [PubMed - indexed for MEDLINE]
-  **12: McClaran SR, Harms CA, Pegelow DF, Dempsey JA.** [Related Articles](#), [Li](#)
-  **Smaller lungs in women affect exercise hyperpnea.**
J Appl Physiol. 1998 Jun;84(6):1872-81.
PMID: 9609779 [PubMed - indexed for MEDLINE]
-  **13: Peslin R, Farre R, Rotger M, Navajas D.** [Related Articles](#), [Li](#)
-  **Effect of expiratory flow limitation on respiratory mechanical impedance: a model study.**
J Appl Physiol. 1996 Dec;81(6):2399-406.
PMID: 9018485 [PubMed - indexed for MEDLINE]
-  **14: Barret J, Chitayat D, Sermer M, Amankwah K, Morrow R, Toi A, Ryan G.** [Related Articles](#), [Li](#)
-  **The prognostic factors in the prenatal diagnosis of the echogenic fetal lung.**
Prenat Diagn. 1995 Sep;15(9):849-53.
PMID: 8559756 [PubMed - indexed for MEDLINE]
-  **15: Yamazaki S.** [Related Articles](#), [Li](#)
-  **Fibrous structure of the joint capsule in the human shoulder.**
Okajimas Folia Anat Jpn. 1990 Aug;67(2-3):127-39.
PMID: 2216306 [PubMed - indexed for MEDLINE]
-  **16: Einarsdottir GH, Stankovich MT, Tu SC.** [Related Articles](#), [Li](#)
-  **Studies of electron-transfer properties of salicylate hydroxylase from Pseudomonas cepacia and effects of salicylate and benzoate binding.**
Biochemistry. 1988 May 3;27(9):3277-85.
PMID: 3390431 [PubMed - indexed for MEDLINE]

Display **Summary** Show: **500** **Sort** **Send to** **Text**

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
Department of Health & Human Services
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books

Search PubMed for ephrin-B3 Go Clear

Limits Preview/Index History Clipboard Details

Display Summary Show: 500 Sort Send to Text

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Items 1 - 41 of 41

One p

1: [Wieland I, Jakubiczka S, Muschke P, Cohen M, Thiele H, Gerlach KL, Adams RH, Wieacker P.](#) Related Articles, Li

Mutations of the ephrin-B1 gene cause craniofrontonasal syndrome. Am J Hum Genet. 2004 Jun;74(6):1209-15. Epub 2004 Apr 29. PMID: 15124102 [PubMed - indexed for MEDLINE]

2: [Bembenek ME, Schmidt S, Li P, Morawiak J, Prack A, Jain S, Roy R, Parsons T, Chee L.](#) Related Articles, Li

Characterization of the kinase domain of the ephrin-B3 receptor tyrosine kinase using a scintillation proximity assay. Assay Drug Dev Technol. 2003 Aug;1(4):555-63. PMID: 15090252 [PubMed - indexed for MEDLINE]

3: [Pascall JC, Brown KD.](#) Related Articles, Li

Intramembrane cleavage of ephrinB3 by the human rhomboid family protease RHBDL2. Biochem Biophys Res Commun. 2004 Apr 23;317(1):244-52. PMID: 15047175 [PubMed - indexed for MEDLINE]

4: [Chen ZY, Sun C, Reuhl K, Bergemann A, Henkemeyer M, Zhou R.](#) Related Articles, Li

Abnormal hippocampal axon bundling in EphB receptor mutant mice. J Neurosci. 2004 Mar 10;24(10):2366-74. PMID: 15014111 [PubMed - indexed for MEDLINE]

5: [Grunwald IC, Korte M, Adelmann G, Plueck A, Kullander K, Adams RH, Frotscher M, Bonhoeffer T, Klein R.](#) Related Articles, Li

Hippocampal plasticity requires postsynaptic ephrinBs. Nat Neurosci. 2004 Jan;7(1):33-40. Epub 2003 Dec 14. PMID: 14699416 [PubMed - indexed for MEDLINE]

6: [Yu G, Luo H, Wu Y, Wu J.](#) Related Articles, Li











Mouse ephrinB3 augments T-cell signaling and responses to T-cell receptor ligation. J Biol Chem. 2003 Nov 21;278(47):47209-16. Epub 2003 Sep 17. PMID: 13679370 [PubMed - indexed for MEDLINE]

7: [Egawa M, Yoshioka S, Higuchi T, Sato Y, Tatsumi K, Fujiwara H, Fujii S.](#) Related Articles, Li


Ephrin B1 is expressed on human luteinizing granulosa cells in corpora lutea of the early luteal phase: the possible involvement of the B class Eph-ephrin system during corpus luteum formation. J Clin Endocrinol Metab. 2003 Sep;88(9):4384-92. PMID: 12970314 [PubMed - indexed for MEDLINE]

8: [Kiehn O, Butt SJ.](#) Related Articles, Li

Physiological, anatomical and genetic identification of CPG neurons in the developing mammalian spinal cord. Prog Neurobiol. 2003 Jul;70(4):347-61. Review. PMID: 12963092 [PubMed - indexed for MEDLINE]

- ☐ **9:** [de Saint-Vis B, Bouchet C, Gautier G, Valladeau J, Caux C, Garrone P.](#) Related Articles, Li
 **Human dendritic cells express neuronal Eph receptor tyrosine kinases: role of EphA2 in regulating adhesion to fibronectin.**
 Blood. 2003 Dec 15;102(13):4431-40. Epub 2003 Aug 07.
 PMID: 12907451 [PubMed - indexed for MEDLINE]
- ☐ **10:** [Tabakoff B, Bhavé SV, Hoffman PL.](#) Related Articles, Li
 **Selective breeding, quantitative trait locus analysis, and gene arrays identify candidate genes for complex drug-related behaviors.**
 J Neurosci. 2003 Jun 1;23(11):4491-8.
 PMID: 12805289 [PubMed - indexed for MEDLINE]
- ☐ **11:** [Brors D, Bodmer D, Pak K, Aletsee C, Schafers M, Dazert S, Ryan AF.](#) Related Articles, Li
 **EphA4 provides repulsive signals to developing cochlear ganglion neurites mediated through ephrin-B2 and -B3.**
 J Comp Neurol. 2003 Jul 14;462(1):90-100.
 PMID: 12761826 [PubMed - indexed for MEDLINE]
- ☐ **12:** [Kubis N, Catala M.](#) Related Articles, Li
 **[Development and maturation of the pyramidal tract]**
 Neurochirurgie. 2003 May;49(2-3 Pt 2):145-53. Review. French.
 PMID: 12746689 [PubMed - indexed for MEDLINE]
- ☐ **13:** [Sanchez-Carbayo M, Belbin TJ, Scotlandi K, Prystowsky M, Baldini N, Childs G, Cordon-Cardo C.](#) Related Articles, Li
 **Expression profiling of osteosarcoma cells transfected with MDR1 and NEO genes: regulation of cell adhesion, apoptosis, and tumor suppression-related genes.**
 Lab Invest. 2003 Apr;83(4):507-17.
 PMID: 12695554 [PubMed - indexed for MEDLINE]
- ☐ **14:** [Howard MA, Rodenas-Ruano A, Henkemeyer M, Martin GK, Lonsbury-Martin BL, Liebl DJ.](#) Related Articles, Li
 **Eph receptor deficiencies lead to altered cochlear function.**
 Hear Res. 2003 Apr;178(1-2):118-30.
 PMID: 12684184 [PubMed - indexed for MEDLINE]
- ☐ **15:** [Pickles JO.](#) Related Articles, Li
 **Expression of Ephs and ephrins in developing mouse inner ear.**
 Hear Res. 2003 Apr;178(1-2):44-51.
 PMID: 12684176 [PubMed - indexed for MEDLINE]
- ☐ **16:** [Kullander K, Butt SJ, Lebreu JM, Lundfald L, Restrepo CE, Rydstrom A, Klein R, Kiehn O.](#) Related Articles, Li
 **Role of EphA4 and EphrinB3 in local neuronal circuits that control walking.**
 Science. 2003 Mar 21;299(5614):1889-92.
 PMID: 12649481 [PubMed - indexed for MEDLINE]
- ☐ **17:** [Bianchi LM, Dinsio K, Davoli K, Gale NW.](#) Related Articles, Li
 **Lac z Histochemistry and immunohistochemistry reveal ephrin-B ligand expression in the inner ear.**
 J Histochem Cytochem. 2002 Dec;50(12):1641-5.
 PMID: 12486086 [PubMed - indexed for MEDLINE]
- ☐ **18:** [Takemoto M, Fukuda T, Sonoda R, Murakami F, Tanaka H, Yamamoto N.](#) Related Articles, Li
 **Ephrin-B3-EphA4 interactions regulate the growth of specific thalamocortical axon populations in vitro.**
 Eur J Neurosci. 2002 Sep;16(6):1168-72.

PMID: 12383247 [PubMed - indexed for MEDLINE]


-  **19:** Varelias A, Koblar SA, Cowled PA, Carter CD, Clayer M. Related Articles, Li



Human osteosarcoma expresses specific ephrin profiles: implications for tumorigenicity and prognosis.

Cancer. 2002 Aug 15;95(4):862-9.

PMID: 12209731 [PubMed - indexed for MEDLINE]


-  **20:** Hindges R, McLaughlin T, Genoud N, Henkemeyer M, O'Leary DD. Related Articles, Li



EphB forward signaling controls directional branch extension and arborization required for dorsal-ventral retinotopic mapping.

Neuron. 2002 Aug 1;35(3):475-87.

PMID: 12165470 [PubMed - indexed for MEDLINE]


-  **21:** Liu W, Ahmad SA, Jung YD, Reinmuth N, Fan F, Bucana CD, Ellis LM. Related Articles, Li



Coexpression of ephrin-Bs and their receptors in colon carcinoma.

Cancer. 2002 Feb 15;94(4):934-9.

PMID: 11920461 [PubMed - indexed for MEDLINE]


-  **22:** Grunwald IC, Korte M, Wolfer D, Wilkinson GA, Unsicker K, Lipp HP, Bonhoeffer T, Klein R. Related Articles, Li



Kinase-independent requirement of EphB2 receptors in hippocampal synaptic plasticity.

Neuron. 2001 Dec 20;32(6):1027-40.

PMID: 11754835 [PubMed - indexed for MEDLINE]


-  **23:** Coonan JR, Greferath U, Messenger J, Hartley L, Murphy M, Boyd AW, Dottori M, Galca MP, Bartlett PF. Related Articles, Li



Development and reorganization of corticospinal projections in EphA4 deficient mice.

J Comp Neurol. 2001 Jul 23;436(2):248-62.

PMID: 11438928 [PubMed - indexed for MEDLINE]


-  **24:** Stuckmann I, Weigmann A, Shevchenko A, Mann M, Huttner WB. Related Articles, Li



Ephrin B1 is expressed on neuroepithelial cells in correlation with neocortical neurogenesis.

J Neurosci. 2001 Apr 15;21(8):2726-37.

PMID: 11306625 [PubMed - indexed for MEDLINE]

-  **25:** Kullander K, Croll SD, Zimmer M, Pan L, McClain J, Hughes V, Zabski S, DeChiara TM, Klein R, Yancopoulos GD, Gale NW. Related Articles, Li



Ephrin-B3 is the midline barrier that prevents corticospinal tract axons from recrossing, allowing for unilateral motor control.

Genes Dev. 2001 Apr 1;15(7):877-88.

PMID: 11297511 [PubMed - indexed for MEDLINE]


-  **26:** Rutherford A. Related Articles, Li



Looking into mirror movement disorder.

Trends Mol Med. 2001 Mar;7(3):101. No abstract available.

PMID: 11286777 [PubMed - indexed for MEDLINE]

-  **27:** Yokoyama N, Romero MI, Cowan CA, Galvan P, Helmbacher F, Charnay P, Parada LF, Henkemeyer M. Related Articles, Li



Forward signaling mediated by ephrin-B3 prevents contralateral corticospinal axons from recrossing the spinal cord midline.

Neuron. 2001 Jan;29(1):85-97.

PMID: 11182083 [PubMed - indexed for MEDLINE]

-  **28:** Tang XX, Zhao H, Robinson ME, Cnaan A, London W, Cohn SL. Related Articles, Li

Cheung NK, Brodeur GM, Evans AE, Ikegaki N.



Prognostic significance of EPHB6, EFNB2, and EFNB3 expressions in neuroblastoma.

Med Pediatr Oncol. 2000 Dec;35(6):656-8.

PMID: 11107140 [PubMed - indexed for MEDLINE]

- ☐ **29:** Tang XX, Zhao H, Robinson ME, Cohen B, Cnaan A, London W, Cohn SL, Cheung NK, Brodeur GM, Evans AE, Ikegaki N. Related Articles, Li



Implications of EPHB6, EFNB2, and EFNB3 expressions in human neuroblastoma.

Proc Natl Acad Sci U S A. 2000 Sep 26;97(20):10936-41.

PMID: 10984508 [PubMed - indexed for MEDLINE]

- ☐ **30:** L'Allemain G.

Related Articles, Li



[Involvement of ephrins and their receptors in oncogenesis]

Bull Cancer. 2000 Jul;87(7-8):529-30. French.

PMID: 10969208 [PubMed - indexed for MEDLINE]

- ☐ **31:** Helbling PM, Saulnier DM, Robinson V, Christiansen JH, Wilkinson DG, Brandli AW. Related Articles, Li



Comparative analysis of embryonic gene expression defines potential interaction sites for Xenopus EphB4 receptors with ephrin-B ligands.

Dev Dyn. 1999 Dec;216(4-5):361-73.

PMID: 10633856 [PubMed - indexed for MEDLINE]

- ☐ **32:** Helbling PM, Saulnier DM, Brandli AW.

Related Articles, Li



The receptor tyrosine kinase EphB4 and ephrin-B ligands restrict angiogenic growth of embryonic veins in Xenopus laevis.

Development. 2000 Jan;127(2):269-78.

PMID: 10603345 [PubMed - indexed for MEDLINE]

- ☐ **33:** van Heumen WR, Claxton C, Pickles JO.

Related Articles, Li



Expression of EphA4 in developing inner ears of the mouse and guinea pig.

Hear Res. 2000 Jan;139(1-2):42-50.

PMID: 10601711 [PubMed - indexed for MEDLINE]

- ☐ **34:** Tang XX, Evans AE, Zhao H, Cnaan A, London W, Cohn SL, Brodeur GM, Ikegaki N. Related Articles, Li



High-level expression of EPHB6, EFNB2, and EFNB3 is associated with low tumor stage and high TrkA expression in human neuroblastomas.

Clin Cancer Res. 1999 Jun;5(6):1491-6.

PMID: 10389937 [PubMed - indexed for MEDLINE]

- ☐ **35:** Miranda JD, White LA, Marcillo AE, Willson CA, Jagid J, Whittmore SR. Related Articles, Li



Induction of Eph B3 after spinal cord injury.

Exp Neurol. 1999 Mar;156(1):218-22.

PMID: 10192794 [PubMed - indexed for MEDLINE]

- ☐ **36:** O'Leary DD, Wilkinson DG.

Related Articles, Li



Eph receptors and ephrins in neural development.

Curr Opin Neurobiol. 1999 Feb;9(1):65-73. Review.

PMID: 10072375 [PubMed - indexed for MEDLINE]

- ☐ **37:** Tang XX, Brodeur GM, Campling BG, Ikegaki N.

Related Articles, Li



Coexpression of transcripts encoding EPHB receptor protein tyrosine kinase and their ephrin-B ligands in human small cell lung carcinoma.

Clin Cancer Res. 1999 Feb;5(2):455-60.

PMID: 10037197 [PubMed - indexed for MEDLINE]

 **38:** [Lin D, Gish GD, Songyang Z, Pawson T.](#)

[Related Articles](#), [Li](#)



The carboxyl terminus of B class ephrins constitutes a PDZ domain binding motif.

J Biol Chem. 1999 Feb 5;274(6):3726-33.

PMID: 9920925 [PubMed - indexed for MEDLINE]

 **39:** [Flanagan JG, Vanderhaeghen P.](#)


[Related Articles](#), [Li](#)



The ephrins and Eph receptors in neural development.

Annu Rev Neurosci. 1998;21:309-45. Review.

PMID: 9530499 [PubMed - indexed for MEDLINE]


 **40:** [Bergemann AD, Zhang L, Chiang MK, Brambilla R, Klein R, Flanagan JG.](#) [Related Articles](#), [Li](#)



Ephrin-B3, a ligand for the receptor EphB3, expressed at the midline of the developing neural tube.

Oncogene. 1998 Jan 29;16(4):471-80.

PMID: 9484836 [PubMed - indexed for MEDLINE]

 **41:** [Tang XX, Pleasure DE, Ikegaki N.](#)




[Related Articles](#), [Li](#)



cDNA cloning, chromosomal localization, and expression pattern of EPLG8, new member of the EPLG gene family encoding ligands of EPH-related protein-tyrosine kinase receptors.

Genomics. 1997 Apr 1;41(1):17-24.

PMID: 9126477 [PubMed - indexed for MEDLINE]

Display **Summary**  Show: **500**  **Sort**  **Send to** **Text**


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show: 20 Sort Send to Text

About Entrez

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

☐ 1: Genomics. 1997 Apr 1;41(1):17-24.

Related Articles, I

FULL-TEXT ARTICLE

cDNA cloning, chromosomal localization, and expression pattern EPLG8, a new member of the EPLG gene family encoding ligand of EPH-related protein-tyrosine kinase receptors.

Tang XX, Pleasure DE, Ikegaki N.

Division of Neurology Research, Children's Hospital of Philadelphia, Pennsylvania 19104-4318, USA.

By screening a human fetal brain cDNA library under low stringency using cDNA encoding the mouse ligand of Cek5 as a probe, we have isolated a novel cDNA belonging to the EPLG gene family. This family encodes ligands of EPH-related tyrosine kinase receptors. Since the novel gene is the eighth member of the EPLG gene family, it is designated EPLG8. The deduced amino acid sequence of EPLG8 suggests that it encodes a transmembrane protein that is more related to those encoded by EPLG2 and EPLG5. We mapped the EPLG8 gene to human chromosome 17p11.2-p13.1 by PCR screening of human-rodent somatic cell hybrid panels. In the midterm fetus, EPLG8 mRNA is expressed at the highest level in brain, followed by heart, kidney, and lung. In the adult, EPLG8 mRNA expression is restricted to brain. These data suggest that LERK-8, the protein encoded by EPLG8, is important in brain development as well as in its maintenance. Moreover, since levels of EPLG8 expression were particularly high in several forebrain subregions compared to other brain subregions, LERK-8 may play a pivotal role in forebrain function.

PMID: 9126477 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show: 20 Sort Send to Text

About Entrez

Text Version

1: Oncogene. 1998 Jan 29;16(4):471-80.

Related Articles, 1

Entrez PubMed


Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central



Ephrin-B3, a ligand for the receptor EphB3, expressed at the midline of the developing neural tube.

Bergemann AD, Zhang L, Chiang MK, Brambilla R, Klein R, Flanagan J

Department of Cell Biology, Harvard Medical School, Boston, Massachusetts 02115, USA.

The ephrins are a family of ligands that bind to Eph family receptor tyrosine kinases, and have been implicated in axon guidance and other patterning processes during vertebrate development. We describe here the identification and characterization of murine ephrin-B3. The cDNA encodes a 340 amino acid transmembrane molecule, most closely related to the two other known transmembrane ligands, ephrin-B1 and ephrin-B2. In addition to homology in their extracellular receptor binding domains, these transmembrane ligands share striking homology between their cytoplasmic domains, with 31 of the last 34 amino acids of ephrin-B3 being identical to ephrin-B2, suggesting functional interactions of the cytoplasmic tail. While most Eph family ligands are promiscuous in their interactions with Eph receptors, binding studies with the receptors known to bind other transmembrane ligands only revealed a high affinity interaction of ephrin-B3 with EphB3, with a dissociation constant of approximately 1 nM. In situ hybridization of mouse embryos showed ephrin-B is expressed prominently at the dorsal and ventral midline of the neural tube, particularly in the floor plate, a structure with key functions in patterning the nervous system. The isolation of this ligand may help to elucidate the molecular basis of patterning activities at the neural tube midline.

PMID: 9484836 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show: 20 Sort Send to Text

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: Exp Neurol. 1999 Mar;156(1):218-22.

Related Articles, I

Full-Text Article

Induction of Eph B3 after spinal cord injury.

Miranda JD, White LA, Marcillo AE, Willson CA, Jagid J, Whittemore S.

Department of Neurological Surgery, University of Miami School of Medicine
1600 Northwest 10th Avenue, R-48, Miami, Florida 33136, USA.

Spinal cord injury (SCI) in adult rats initiates a cascade of events producing a nonpermissive environment for axonal regeneration. This nonfavorable environment could be due to the expression of repulsive factors. The Eph receptor protein tyrosine kinases and their respective ligands (ephrins) are families of molecules that play a major role in axonal pathfinding and target recognition during central nervous system (CNS) development. Their mechanism of action mediated by repellent forces between receptor and ligand. The possible role that these molecules play after CNS trauma is unknown. We hypothesized that an increase in the expression of Eph proteins and/or ephrins may be one of the molecular cues that restrict axonal regeneration after SCI. Rats received a contusive SCI at T10 and in situ hybridization studies 7 days posttrauma demonstrated: (i) a marked up-regulation of Eph B3 mRNA in cells located in white matter at the lesion epicenter, but not rostral or caudal to the injury site, (ii) an increase in Eph B3 mRNA in neurons in the ventral horn and intermediate zone of the gray matter, rostral and caudal to the lesion. Immunohistochemical analyses localizing Eph B3 protein were consistent with the mRNA results. Colocalization studies performed in injured animals demonstrated increased Eph B3 expression in white matter astrocytes and motor neurons of the gray matter. These results suggest that Eph B3 may contribute to the unfavorable environment for axonal regeneration after SCI. Copyright 1999 Academic Press.

PMID: 10192794 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show: 20 Sort Send to Text

About Entrez

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

☐ 1: Genes Dev. 2001 Apr 1;15(7):877-88.

Related Articles, ...

FREE full text article of
www.genesdev.org

Ephrin-B3 is the midline barrier that prevents corticospinal tract axons from recrossing, allowing for unilateral motor control.

Kullander K, Croll SD, Zimmer M, Pan L, McClain J, Hughes V, Zabiski DeChiara TM, Klein R, Yancopoulos GD, Gale NW.


European Molecular Biology Laboratory, D-69117 Heidelberg, Germany.

Growing axons follow highly stereotypical pathways, guided by a variety of attractive and repulsive cues, before establishing specific connections with distal targets. A particularly well-known example that illustrates the complexity of axonal migration pathways involves the axonal projections of motor neurons located in the motor cortex. These projections take a complex route during which they first cross the midline, then form the corticospinal tract, and ultimately connect with motor neurons in the contralateral side of the spinal cord. These obligatory contralateral connections account for why one side of the brain controls movement on the opposing side of the body. The netrins and slits provide well-known midline signals that regulate axonal crossings at the midline. Here we report that a member of the ephrin family, ephrin-B3, also plays a key role at the midline to regulate axonal crossing. In particular, we show that ephrin-B3 acts as the midline barrier that prevents corticospinal tract projections from recrossing when they enter the spinal gray matter. We report that in ephrin-B3(-/-) mice, corticospinal tract projections freely recross in the spinal gray matter, such that the motor cortex on one side of the brain now provides bilateral input to the spinal cord. This neuroanatomical abnormality in ephrin-B3(-/-) mice correlates with loss of unilateral motor control, yielding mice that simultaneously move their right and left limbs and thus have a peculiar hopping gait quite unlike the alternate step gait displayed by normal mice. The corticospinal and walking defects in ephrin-B3(-/-) mice resemble those recently reported for mice lacking the EphA4 receptor, which binds ephrin-B3 as well as other ephrins, suggesting that the binding of EphA4-bearing axonal processes to ephrin-B3 at the midline provides the repulsive signal that prevents corticospinal tract projections from recrossing the midline in the developing spinal cord.

PMID: 11297511 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search for

Limits Preview/Index History Clipboard Details

Show:

About Entrez

Text Version

☐ 1: Eur J Neurosci. 2002 Sep;16(6):1168-72.

Related Articles, 1

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central



Ephrin-B3-EphA4 interactions regulate the growth of specific thalamocortical axon populations in vitro.

Takemoto M, Fukuda T, Sonoda R, Murakami F, Tanaka H, Yamamoto I

Neuroscience Laboratories, Graduate School of Frontier Biosciences, Osaka University, Toyonaka, Osaka 560-8531, Japan.

The role was studied of ephrin-B3, a ligand of the Eph family of tyrosine kinase receptors, in the formation of cortical connectivity. In situ hybridization and immunohistochemistry showed that EphA4, a receptor of ephrin-B3, was expressed in the lateral thalamus (visual and somatosensory thalamus) of the developing rat brain, but not in the medial thalamic nuclei which project to the limbic cortex. Correspondingly, ephrin-B3 was expressed strongly in the developing limbic cortex including amygdala, entorhinal cortex and hippocampus. To examine the action of ephrin-B3 on thalamic axons, either lateral or medial thalamic explants were cultured on membranes obtained from ephrin-B3-expressing COS cells. Axonal growth was inhibited for cells from the lateral thalamus but not from the medial thalamus. These results suggest that ephrin-B3 contributes to regional specificity by suppressing axonal growth of lateral thalamic neurons.

PMID: 12383247 [PubMed - indexed for MEDLINE]

Show:


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search for

Limits Preview/Index History Clipboard Details

Show: Sort

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: J Biol Chem. 2003 Nov 21;278(47):47209-16. Epub 2003 Sep 17.

Related Arti

FREE full text article at
www.jbc.org

Mouse ephrinB3 augments T-cell signaling and responses to T-cell receptor ligation.

Yu G, Luo H, Wu Y, Wu J.

Laboratory of Immunology and the Nephrology Service of Notre Dame Hospital Centre Hospitalier de l'Universite de Montreal, Universite de Montreal, Montreal, Quebec H2L 4M1, Canada.

Ephrins (EFN) are cell-surface ligands of Ephs, the largest family of cell-surface receptor tyrosine kinases. The function of EFNs in the immune system has not been well studied, although some EFNs and Ephs are expressed at high levels in certain leukocytes. We report here that EFN3 and its receptors (collectively called EFN3Rs, as EFN3 binds to multiple EphBs) were expressed in peripheral T cells and monocytes/macrophages, with T cells being the dominant EFN3+ and EFN3R+ cell type. Solid-phase EFN3-Fc in the presence of suboptimal anti-CD3 crosslinking enhanced T-cell responses in terms of proliferation, activation marker expression, interferon-gamma but not interleukin-2 production, and cytotoxic T-cell activity. EFN3R costimulation in the presence of phorbol 12-myristate 13-acetate was insensitive to cyclosporin A, similar to CD28 costimulation, suggesting they might share a part of the signaling pathway. After crosslinking, T-cell receptor and EFN3R congregated into aggregated rafts, and this provided a morphological basis for signaling pathway of T-cell receptor and EFN3R to interact. Solid-phase EFN3-Fc augmented p38 and p44/42 MAPK activation further downstream of the signaling pathway. These data suggest that EFN3 is important in T-cell/T-cell and T-cell/antigen-presenting cell collaboration to enhance T-cell activation and function.

PMID: 13679370 [PubMed - indexed for MEDLINE]

Show: Sort


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09



Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show: 20 Sort Send to Text

About Entrez

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services


Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

☐ 1: J Comp Neurol. 2003 Jul 14;462(1):90-100.

Related Articles, i



EphA4 provides repulsive signals to developing cochlear ganglion neurites mediated through ephrin-B2 and -B3.

Brors D, Bodmer D, Pak K, Aletsee C, Schafers M, Dazert S, Ryan AF.

Department of Surgery, Division Otolaryngology and Neurosciences, University of California, San Diego School of Medicine and Veterans Administration Medical Center, La Jolla 92093, USA.

The ephrins and Eph receptors make up two large families of bi-directional signaling molecules that are known to play a role in the development of the nervous system. Recently, expression of EphA4 in the developing cochlea was shown, with strong expression in cells lining the osseous spiral lamina (OSL) through which afferent dendrites must pass to reach the organ of Corti (OC). It was also demonstrated that ephrin-B2 and -B3, both of which are known to interact with EphA4, are expressed by spiral ganglion (SG) neurons. To investigate the functional role of EphA4 in the development of inner ear neuro neonatal rat SG explants were cultured for 72 hours on uniformly coated surface or near stripes of EphA4/IgG-Fc-chimera. Control explants were cultured on or near IgG-Fc and EphA1/IgG-Fc-chimera. To assess the roles of ephrin-B2 and B3 in EphA4 signaling, SG explants were cultured with or without anti-ephrin and/or -B3 blocking antibodies. Growth patterns of SG neurites at the border of EphA4 receptor stripes showed repulsion, characterized by turning, stopping and/or reversal. In the case of IgG-Fc and EphA1, the neurites grew straight over the stripes. Treatment with either anti-ephrin-B2 or -B3 blocking antibodies significantly reduced the repulsive effect of an EphA4 stripe. Moreover, when both antibodies were used together, neurites crossed onto EphA4 stripes with no evidence of repulsion. The results suggest that EphA4 provides repulsive signals to SG neurites in the developing cochlea, and that ephrin-B2 and -B3 together mediate this response. Copyright 2003 Wiley-Liss, Inc.

PMID: 12761826 [PubMed - indexed for MEDLINE]


Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)



Entrez PubMed Nucleotide Protein Genomes Structure OMIM PMC Journals Books

Search PubMed for ephrin AND vascularization Go Clear

Limits Preview/Index History Clipboard Details

Display Summary Show: 500 Sort Send to Text

About Entrez

Items 1 - 35 of 35

One page

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Umeda N, Ozaki H, Hayashi H, Oshima K.

Related Articles, Li



Expression of ephrinB2 and its receptors on fibroproliferative membranes in ocular angiogenic diseases.

Am J Ophthalmol. 2004 Aug;138(2):270-9.

PMID: 15289137 [PubMed - indexed for MEDLINE]

2: Liu W, Jung YD, Ahmad SA, McCarty MF, Stoeltzing O, Reinmuth N, Fan F, Ellis LM.

Related Articles, Li



Effects of overexpression of ephrin-B2 on tumour growth in human colorectal cancer.

Br J Cancer. 2004 Apr 19;90(8):1620-6.

PMID: 15083195 [PubMed - indexed for MEDLINE]

3: Noren NK, Lu M, Freeman AL, Koolpe M, Pasquale EB.

Related Articles, Li



Interplay between EphB4 on tumor cells and vascular ephrin-B2 regulates tumor growth.

Proc Natl Acad Sci U S A. 2004 Apr 13;101(15):5583-8. Epub 2004 Apr 05.

PMID: 15067119 [PubMed - indexed for MEDLINE]

4: Patan S.

Related Articles, Li



Vasculogenesis and angiogenesis.

Cancer Treat Res. 2004;117:3-32. Review.

PMID: 15015550 [PubMed - indexed for MEDLINE]

5: Huang J, Soffer SZ, Kim ES, McCrudden KW, Huang J, New T, Manley CA, Middlesworth W, O'Toole K, Yamashiro DJ, Kandel JJ.

Related Articles, Li



Vascular remodeling marks tumors that recur during chronic suppression of angiogenesis.

Mol Cancer Res. 2004 Jan;2(1):36-42.

PMID: 14757844 [PubMed - indexed for MEDLINE]

6: Bicknell R, Harris AL.

Related Articles, Li



Novel angiogenic signaling pathways and vascular targets.

Annu Rev Pharmacol Toxicol. 2004;44:219-38. Review.

PMID: 14744245 [PubMed - indexed for MEDLINE]

7: Sawai Y, Tamura S, Fukui K, Ito N, Imanaka K, Sacki A, Sakuda S, Kiso S, Matsuzawa Y.

Related Articles, Li



Expression of ephrin-B1 in hepatocellular carcinoma: possible involvement in neovascularization.

J Hepatol. 2003 Dec;39(6):991-6.

PMID: 14642617 [PubMed - indexed for MEDLINE]

8: Shibuya M.


Related Articles, Li




VEGF-receptor inhibitors for anti-angiogenesis.

Nippon Yakurigaku Zasshi. 2003 Dec;122(6):498-503. Review.


PMID: 14639004 [PubMed - indexed for MEDLINE]

 **9:** [Augustin HG, Reiss Y.](#)


[Related Articles](#), [Li](#)

 **EphB receptors and ephrinB ligands: regulators of vascular assembly and homeostasis.**


Cell Tissue Res. 2003 Oct;314(1):25-31. Epub 2003 Aug 02. Review.
PMID: 12905065 [PubMed - indexed for MEDLINE]

 **10:** [Sullivan DC, Bicknell R.](#)


[Related Articles](#), [Li](#)

 **New molecular pathways in angiogenesis.**


Br J Cancer. 2003 Jul 21;89(2):228-31. Review.
PMID: 12865906 [PubMed - indexed for MEDLINE]

 **11:** [Andres AC, Munarini N, Djonov V, Bruneau S, Zuercher G, Loercher S, Rohrbach V, Ziemiecki A.](#)


[Related Articles](#), [Li](#)

 **EphB4 receptor tyrosine kinase transgenic mice develop glomerulopathies reminiscent of aglomerular vascular shunts.**

Mech Dev. 2003 Apr;120(4):511-6.
PMID: 12676328 [PubMed - indexed for MEDLINE]

 **12:** [Marme D.](#)


[Related Articles](#), [Li](#)

 **VEGFs, angiopoietins, Ephrins and their receptors: putative targets for tumor therapy?**


Ann Hematol. 2002;81 Suppl 2:S66. No abstract available.
PMID: 12611080 [PubMed - indexed for MEDLINE]

 **13:** [Augustin HG.](#)


[Related Articles](#), [Li](#)

 **[Angiogenesis research--quo vadis?]**


Ophthalmologe. 2003 Feb;100(2):104-10. Review. German.
PMID: 12589453 [PubMed - indexed for MEDLINE]

 **14:** [Takahashi T, Takahashi K, St John PL, Fleming PA, Tomemori T, Watanabe T, Abrahamson DR, Drake CJ, Shirasawa T, Daniel TO.](#)


[Related Articles](#), [Li](#)

 **A mutant receptor tyrosine phosphatase, CD148, causes defects in vascular development.**


Mol Cell Biol. 2003 Mar;23(5):1817-31.
PMID: 12588999 [PubMed - indexed for MEDLINE]

 **15:** [Straume O, Jackson DG, Akslen LA.](#)


[Related Articles](#), [Li](#)

 **Independent prognostic impact of lymphatic vessel density and presence of low-grade lymphangiogenesis in cutaneous melanoma.**

Clin Cancer Res. 2003 Jan;9(1):250-6.
PMID: 12538477 [PubMed - indexed for MEDLINE]

 **16:** [Kaban K, Herbst RS.](#)


[Related Articles](#), [Li](#)

 **Angiogenesis as a target for cancer therapy.**


Hematol Oncol Clin North Am. 2002 Oct;16(5):1125-71. Review.
PMID: 12512387 [PubMed - indexed for MEDLINE]

 **17:** [Meyer S, Hafner C, Vogt T.](#)


[Related Articles](#), [Li](#)

 **[Role of receptor tyrosine kinase in the angiogenesis]**


Hautarzt. 2002 Sep;53(9):629-42. Review. German. No abstract available.
PMID: 12432901 [PubMed - indexed for MEDLINE]

 **18:** [Brantley DM, Cheng N, Thompson EJ, Lin Q, Brekken RA, Thorpe PE, Muraoka RS, Cerretti DP, Pozzi A, Jackson D, Lin C, Chen J.](#)




















[Related Articles](#), [Li](#)

 **Soluble Eph A receptors inhibit tumor angiogenesis and progression in vivo.**


Oncogene. 2002 Oct 10;21(46):7011-26.
PMID: 12370823 [PubMed - indexed for MEDLINE]

 **19:** [Papoutsis M, Othman-Hassan K, Christ B, Patel K, Wilting J.](#)

[Related Articles](#), [Li](#)


-  **Development of an arterial tree in C6 gliomas but not in A375 melanomas.**
Histochem Cell Biol. 2002 Sep;118(3):241-9. Epub 2002 Jul 13.
PMID: 12271360 [PubMed - indexed for MEDLINE]
-  **20:** [Nakamoto M, Bergemann AD.](#) Related Articles, Li
-  **Diverse roles for the Eph family of receptor tyrosine kinases in carcinogenes**
Microsc Res Tech. 2002 Oct 1;59(1):58-67. Review.
PMID: 12242697 [PubMed - indexed for MEDLINE]
-  **21:** [Steinle JJ, Meininger CJ, Forough R, Wu G, Wu MH, Granger HJ.](#) Related Articles, Li
-  **Eph B4 receptor signaling mediates endothelial cell migration and proliferation via the phosphatidylinositol 3-kinase pathway.**
J Biol Chem. 2002 Nov 15;277(46):43830-5. Epub 2002 Sep 13.
PMID: 12235151 [PubMed - indexed for MEDLINE]
-  **22:** [Straume O, Akslen LA.](#) Related Articles, Li
-  **Importance of vascular phenotype by basic fibroblast growth factor, and influence of the angiogenic factors basic fibroblast growth factor/fibroblast growth factor receptor-1 and ephrin-A1/EphA2 on melanoma progression.**
Am J Pathol. 2002 Mar;160(3):1009-19.
PMID: 11891198 [PubMed - indexed for MEDLINE]
-  **23:** [Cheng N, Brantley DM, Chen J.](#) Related Articles, Li
-  **The ephrins and Eph receptors in angiogenesis.**
Cytokine Growth Factor Rev. 2002 Feb;13(1):75-85. Review.
PMID: 11750881 [PubMed - indexed for MEDLINE]
-  **24:** [Kuwano M, Fukushi J, Okamoto M, Nishie A, Goto H, Ishibashi T, Ono M.](#) Related Articles, Li
-  **Angiogenesis factors.**
Intern Med. 2001 Jul;40(7):565-72. Review.
PMID: 11506294 [PubMed - indexed for MEDLINE]
-  **25:** [Gale NW, Baluk P, Pan L, Kwan M, Holash J, DeChiara TM, McDonald DM, Yancopoulos GD.](#) Related Articles, Li
-  **Ephrin-B2 selectively marks arterial vessels and neovascularization sites in the adult, with expression in both endothelial and smooth-muscle cells.**
Dev Biol. 2001 Feb 15;230(2):151-60.
PMID: 11161569 [PubMed - indexed for MEDLINE]
-  **26:** [Shin D, Garcia-Cardena G, Hayashi S, Gerety S, Asahara T, Stavrakis G, Isner J, Folkman J, Gimbrone MA Jr, Anderson DJ.](#) Related Articles, Li
-  **Expression of ephrinB2 identifies a stable genetic difference between arterial and venous vascular smooth muscle as well as endothelial cells, and marks subsets of microvessels at sites of adult neovascularization.**
Dev Biol. 2001 Feb 15;230(2):139-50.
PMID: 11161568 [PubMed - indexed for MEDLINE]
-  **27:** [Ogawa K, Pasqualini R, Lindberg RA, Kain R, Freeman AL, Pasquale EB.](#) Related Articles, Li
-  **The ephrin-A1 ligand and its receptor, EphA2, are expressed during tumor neovascularization.**
Oncogene. 2000 Dec 7;19(52):6043-52.
PMID: 11146556 [PubMed - indexed for MEDLINE]
-  **28:** [Umess LD, Sorensen LK, Li DY.](#) Related Articles, Li
-  **Arteriovenous malformations in mice lacking activin receptor-like kinase-1.**
Nat Genet. 2000 Nov;26(3):328-31.
PMID: 11062473 [PubMed - indexed for MEDLINE]

☐ **29:** [Yancopoulos GD, Davis S, Gale NW, Rudge JS, Wiegand SJ, Holash J.](#) Related Articles, LI

 **Vascular-specific growth factors and blood vessel formation.**
Nature. 2000 Sep 14;407(6801):242-8. Review.
PMID: 11001067 [PubMed - indexed for MEDLINE]


☐ **30:** [Yuan K, Jin YT, Lin MT.](#)

Related Articles, LI


 **Expression of Tie-2, angiopoietin-1, angiopoietin-2, ephrinB2 and EphB4 in pyogenic granuloma of human gingiva implicates their roles in inflammatory angiogenesis.**
J Periodontal Res. 2000 Jun;35(3):165-71.
PMID: 10929871 [PubMed - indexed for MEDLINE]

☐ **31:** [Myers C, Charboneau A, Boudreau N.](#)


Related Articles, LI

 **Homeobox B3 promotes capillary morphogenesis and angiogenesis.**
J Cell Biol. 2000 Jan 24;148(2):343-51.
PMID: 10648567 [PubMed - indexed for MEDLINE]

☐ **32:** [Easty DJ, Hill SP, Hsu MY, Fallowfield ME, Florenes VA, Herlyn M, Bennett DC.](#) Related Articles, LI


 **Up-regulation of ephrin-A1 during melanoma progression.**
Int J Cancer. 1999 Oct 22;84(5):494-501.
PMID: 10502726 [PubMed - indexed for MEDLINE]

☐ **33:** [Adams RH, Wilkinson GA, Weiss C, Diella F, Gale NW, Deutsch U, Risau W, Klein R.](#) Related Articles, LI

 **Roles of ephrinB ligands and EphB receptors in cardiovascular development demarcation of arterial/venous domains, vascular morphogenesis, and sprouting angiogenesis.**
Genes Dev. 1999 Feb 1;13(3):295-306.
PMID: 9990854 [PubMed - indexed for MEDLINE]


☐ **34:** [Chiarugi V, Magnelli L, Gallo O.](#)

Related Articles, LI

 **Cox-2, iNOS and p53 as play-makers of tumor angiogenesis (review).**
Int J Mol Med. 1998 Dec;2(6):715-9. Review.
PMID: 9850741 [PubMed - indexed for MEDLINE]

☐ **35:** [Pandey A, Shao H, Marks RM, Polverini PJ, Dixit VM.](#)

Related Articles, LI

 **Role of B61, the ligand for the Eck receptor tyrosine kinase, in TNF-alpha-induced angiogenesis.**
Science. 1995 Apr 28;268(5210):567-9.
PMID: 7536959 [PubMed - indexed for MEDLINE]

Display Show:

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 13 2004 06:44:09